

# OBSERVATIONS & RECOMMENDATIONS

After reviewing data collected from **CENTER POND, STODDARD**, the program coordinators have made the following observations and recommendations:

We would like to thank your group for sampling your pond **once** this summer. However, we encourage your monitoring group to sample **additional** times each summer. Typically we recommend that monitoring groups sample **three times** per summer (once in **June, July**, and **August**). We understand that the number of sampling events you decide to conduct per summer will depend upon volunteer availability, and your monitoring group's water monitoring goals and funding availability. However, with a limited amount of data it is difficult to determine accurate and representative water quality trends. Since weather patterns and activity in the watershed can change throughout the summer, from year to year, and even from hour to hour during a rain event, it is a good idea to sample the pond at least once per month over the course of the season.

If you are having difficulty finding volunteers to help sample, or to pick-up or drop-off equipment at one of the laboratories, please give the VLAP Coordinator a call and we will try to help you work out an arrangement.

We would like to encourage your monitoring group to formally participate in the DES Weed Watchers program, a volunteer program dedicated to monitoring the lakes and ponds for the presence of exotic aquatic plants. This program only involves a small amount of time during the summer months. Volunteers survey their waterbody once a month from **June** through **September**. To survey, volunteers slowly boat, or even snorkel, around the perimeter of the waterbody and any islands it may contain. Using the materials provided in the Weed Watchers Kit, volunteers look for any species that are of suspicion. After a trip or two around the waterbody, volunteers will have a good knowledge of its plant community and will immediately notice even the most subtle changes. If a suspicious plant is found, the volunteers will send a specimen to DES for identification. If the plant specimen is an exotic, a biologist will visit the site to determine the extent of the problem.

and to formulate a plan of action to control the nuisance infestation. Remember that early detection is the key to controlling the spread of exotic plants.

If you would like to help protect your lake or pond from exotic plants, contact Amy Smagula, Exotic Species Program Coordinator, at 271-2248 or visit the Weed Watchers web page at [www.des.state.nh.us/wmb/exoticspecies/survey.htm](http://www.des.state.nh.us/wmb/exoticspecies/survey.htm).

#### **FIGURE INTERPRETATION**

- **Figure 1 and Table 1:** The graphs in Figure 1 (Appendix A) show the historical and current year chlorophyll-a concentration in the water column. Table 1 (Appendix B) lists the maximum, minimum, and mean concentration for each sampling season that the pond has been monitored through the program.

Chlorophyll-a, a pigment found in plants, is an indicator of the algal abundance. Because algae are usually microscopic plants that contain chlorophyll-a, and are naturally found in lake ecosystems, the chlorophyll-a concentration measured in the water gives an estimation of the algal concentration or lake productivity. **The median summer chlorophyll-a concentration for New Hampshire's lakes and ponds is 4.58 mg/m<sup>3</sup>.**

The current year data (the top graph) show that the chlorophyll-a concentration in **July** was ***slightly greater than*** the state median and was ***approximately equal to*** the similar lake median (refer to Appendix F for more information about the similar lake median).

Overall, visual inspection of the historical data trend line (the bottom graph) shows a ***variable*** in-lake chlorophyll-a trend since monitoring began. Specifically, the mean concentration has ***fluctuated between approximately 3.5 and 11.8 mg/m<sup>3</sup> since 1988.***

In the 2006 annual report, since your group will have sampled the chlorophyll-a concentration at the deep spot for at least 10 consecutive years, we will conduct a statistical analysis of the historic data to determine if there has been a significant change in the annual mean since monitoring began.

While algae are naturally present in all lakes/ponds, an excessive or increasing amount of any type is not welcomed. In freshwater lakes/ponds, phosphorus is the nutrient that algae depend upon for growth. Algal concentrations may increase with an increase in nonpoint sources of phosphorus loading from the watershed, or in-

lake sources of phosphorus loading (such as phosphorus releases from the sediments). Therefore, it is extremely important for volunteer monitors to continually educate residents about how activities within the watershed can affect phosphorus loading and pond quality.

- **Figure 2 and Table 3:** The graphs in Figure 2 (Appendix A) show historical and current year data for pond transparency. Table 3 (Appendix B) lists the maximum, minimum and mean transparency data for each sampling season that the pond has been monitored through the program.

Volunteer monitors use the Secchi-disk, a 20 cm disk with alternating black and white quadrants, to measure water clarity (how far a person can see into the water). Transparency, a measure of water clarity, can be affected by the amount of algae and sediment from erosion, as well as the natural colors of the water. **The median summer transparency for New Hampshire's lakes and ponds is 3.2 meters.**

The current year data (the top graph) show that the in-lake transparency in **July** was *slightly less than* the state median and *slightly greater than* the similar lake median (refer to Appendix F for more information about the similar lake median).

Overall, visual inspection of the historical data trend line (the bottom graph) shows a *variable, but overall increasing (meaning improving)*, transparency trend since monitoring began. Specifically, the mean transparency has *fluctuated between approximately 2.5 and 4.9 meters* since 1988.

As previously discussed, since your group will have sampled the transparency at the deep spot for at least 10 consecutive years, the 2006 annual report will include a statistical analysis of the historic data to determine if there has been a significant change in the annual mean since monitoring began.

Typically, high intensity rainfall causes erosion of sediments into lakes/ponds and streams, thus decreasing clarity. Efforts should continually be made to stabilize stream banks, pond shorelines, disturbed soils within the watershed, and especially dirt roads located immediately adjacent to the edge of tributaries and the pond. Guides to Best Management Practices designed to reduce, and possibly even eliminate, nonpoint source pollutants, such as sediment loading, are available from DES upon request.

- **Figure 3 and Table 8:** The graphs in Figure 3 (Appendix A) show the amount of phosphorus in the epilimnion (the upper layer) and the hypolimnion (the lower layer); the inset graphs show current year data. Table 8 (Appendix B) lists the annual maximum, minimum, and median concentration for each deep spot layer and each tributary since the pond has joined the program.

Phosphorus is the limiting nutrient for plant and algae growth in New Hampshire's freshwater lakes and ponds. Too much phosphorus in a pond can lead to increases in plant and algal growth over time. **The median summer total phosphorus concentration in the epilimnion (upper layer) of New Hampshire's lakes and ponds is 12 ug/L. The median summer phosphorus concentration in the hypolimnion (lower layer) is 14 ug/L.**

The current year data for the epilimnion show that the phosphorus concentration on the **July** sampling event was *slightly less than* the state median and similar lake median.

The current year data for the hypolimnion show that the phosphorus concentration on the **July** sampling event was *greater than* the state median and similar lake median.

The turbidity of the hypolimnion (lower layer) sample was *slightly elevated (3.47 NTUs)* on the **July** sampling event. The turbidity in the hypolimnetic sample has ranged between approximately **0.94** and **5.64 NTUs** since monitoring began which suggests that the lake bottom is covered by a thick organic layer of sediment which is easily disturbed. When the pond bottom is disturbed, sediment, which typically contains attached phosphorus, is released into the water column. When collecting the hypolimnion sample, make sure that there is no sediment in the Kemmerer Bottle before filling the sample bottles.

Overall, visual inspection of the historical data trend line for the epilimnion and hypolimnion shows an overall *decreasing (meaning improving)* phosphorus trend since monitoring began in **1988**.

As previously discussed, since your group will have sampled the phosphorus concentration at the deep spot for at least 10 consecutive years, the 2006 annual report will include a statistical analysis of the historic data to determine if there has been a significant change in the annual mean since monitoring began.

One of the most important approaches to reducing phosphorus loading to a waterbody is to continually educate watershed residents about its sources and how excessive amounts can adversely impact

the ecology and value of lakes and ponds. Phosphorus sources within a lake or pond's watershed typically include septic systems, animal waste, lawn fertilizer, road and construction erosion, and natural wetlands.

#### **TABLE INTERPRETATION**

➤ **Table 2: Phytoplankton**

Table 2 (Appendix B) lists the current and historical phytoplankton species observed in the pond. Specifically, this table lists the three most dominant phytoplankton species observed in the sample and their relative abundance in the sample.

The dominant phytoplankton species observed in the **July** phytoplankton sample were ***Tabellaria* (diatom)**, ***Asterionella* (diatom)**, and ***Synura* (golden-brown)**.

Phytoplankton populations undergo a natural succession during the growing season (Please refer to the “Biological Monitoring Parameters” section of this report for a more detailed explanation regarding seasonal plankton succession). Diatoms and golden-brown algae are typical in New Hampshire's less productive lakes and ponds.

➤ **Table 4: pH**

Table 4 (Appendix B) presents the in-lake and tributary current year and historical pH data.

pH is measured on a logarithmic scale of 0 (acidic) to 14 (basic). pH is important to the survival and reproduction of fish and other aquatic life. A pH below 6.0 limits the growth and reproduction of fish. A pH between 6.0 and 7.0 is ideal for fish. The median pH value for the epilimnion (upper layer) in New Hampshire's lakes and ponds is **6.6**, which indicates that the surface waters in the state are slightly acidic. For a more detailed explanation regarding pH, please refer to the “Chemical Monitoring Parameters” section of this report.

The pH at the deep spot on the **July** sampling event ranged from **5.47** in the hypolimnion to **5.76** in the epilimnion, which means that the water is **acidic**.

It is important to point out that the pH in the hypolimnion (lower layer) was **lower (more acidic)** than in the epilimnion (upper layer). This increase in acidity near the lake bottom is likely due the decomposition of organic matter and the release of acidic by-products into the water column.

Due to the presence of granite bedrock in the state and acid deposition (from snowmelt, rainfall, and atmospheric particulates) in New Hampshire, there is not much that can be done to effectively increase pond pH.

➤ **Table 5: Acid Neutralizing Capacity**

Table 5 (Appendix B) presents the current year and historical epilimnetic ANC for each year the pond has been monitored through VLAP.

Buffering capacity (ANC) describes the ability of a solution to resist changes in pH by neutralizing the acidic input. The median ANC value for New Hampshire's lakes and ponds is **4.9 mg/L**, which indicates that many lakes and ponds in the state are at least "moderately vulnerable" to acidic inputs. For a more detailed explanation, please refer to the "Chemical Monitoring Parameters" section of this report.

The mean Acid Neutralizing Capacity (ANC) of the epilimnion (the upper layer) on the **September** sampling event was **1.3 mg/L** this season, which is ***much less than*** the state median. In addition, this indicates that the pond is ***extremely vulnerable*** to acidic inputs (such as acid precipitation).

➤ **Table 6: Conductivity**

Table 6 (Appendix B) presents the current and historical conductivity values for tributaries and in-lake data. Conductivity is the numerical expression of the ability of water to carry an electric current (which is determined by the number of negatively charged ions from metals, salts, and minerals in the water column). The median conductivity value for New Hampshire's lakes and ponds is **40.0 uMhos/cm**. For a more detailed explanation, please refer to the "Chemical Monitoring Parameters" section of this report.

The conductivity in the epilimnion on the July sampling event at the deep spot this season was **16.83 uMhos/cm**, which is ***much less than*** state median.

The conductivity in the pond is relatively ***stable*** and ***low***. Typically conductivity levels greater than 100 uMhos/cm indicate the influence of human activities on surface water quality. These activities include septic system leachate, agricultural runoff, iron deposits, and road runoff (which contains road salt during the spring snow melt). The low conductivity level in the pond is an indication of the low amount of pollutants and erosion in the watershed. We hope this trend

continues!

➤ **Table 8: Total Phosphorus**

Table 8 (Appendix B) presents the current year and historical total phosphorus data for in-lake and tributary stations. Phosphorus is the nutrient that limits the algae's ability to grow and reproduce. Please refer to the "Chemical Monitoring Parameters" section of this report for a more detailed explanation.

The phosphorus concentration in the **Inlet** was **relatively low** this season, which is good news. However, we recommend that your monitoring group sample the major tributaries to the pond soon after snow-melt and periodically during rain storms to determine if the phosphorus concentration is **elevated** in the tributaries during these times. Typically, the majority of nutrient loading to a pond occurs in the spring during snowmelt and during intense rain storms that cause surface runoff and erosion within the watershed.

*For a detailed explanation on how to conduct rain event sampling please refer to the 2002 VLAP Annual Report "Special Topic Article" or contact the VLAP Coordinator.*

➤ **Table 9 and Table 10: Dissolved Oxygen and Temperature Data**

Table 9 (Appendix B) shows the dissolved oxygen/temperature profile(s) for the 2005 sampling season. Table 10 (Appendix B) shows the historical and current year dissolved oxygen concentration in the hypolimnion (lower layer). The presence of dissolved oxygen is vital to fish and amphibians in the water column and also to bottom-dwelling organisms. Please refer to the "Chemical Monitoring Parameters" section of this report for a more detailed explanation.

The dissolved oxygen concentration was **lower in the hypolimnion (lower layer) than in the epilimnion (upper layer)** at the deep spot of the pond on the **July** sampling event. As stratified ponds age, and as the summer progresses, oxygen typically becomes **depleted** in the hypolimnion by the process of decomposition. Specifically, the loss of oxygen in the hypolimnion results primarily from the process of biological breakdown of organic matter (i.e.; biological organisms use oxygen to break down organic matter), both in the water column and particularly at the bottom of the pond where the water meets the sediment. When oxygen levels are depleted to less than 1 mg/L in the hypolimnion (as it has been during previous annual biologist visits), the phosphorus that is normally bound up in the sediment may be re-released into the water column (a process referred to as **internal phosphorus loading**).

➤ **Table 11: Turbidity**

Table 11 (Appendix B) lists the current year and historical data for in-lake and tributary turbidity. Turbidity in the water is caused by suspended matter, such as clay, silt, and algae. Water clarity is strongly influenced by turbidity. Please refer to the “Other Monitoring Parameters” section of this report for a more detailed explanation.

The **End Camp Brook** turbidity continued to be **relatively low** on the **July** sampling event which suggests that erosion may not be a major contributor of sediment and nutrient loading to the pond in this area of the watershed.

However, on the annual biologist visit in **July**, a sediment delta was observed in the beach/launch area to the lake. Erosion was observed along the dirt/gravel road in this area. It appeared that land located across the road had recent been cleared.

We recommend that your monitoring group conduct a stream survey and storm event sampling in this area of this watershed. This additional sampling may allow us to determine what is causing the **elevated** levels of turbidity.

*For a detailed explanation on how to conduct rain event sampling and stream surveys, please refer to the 2002 VLAP Annual Report “Special Topic Article” or contact the VLAP Coordinator.*

➤ **Table 12: Bacteria (*E.coli*)**

Table 12 lists the current year and historical data for bacteria (*E.coli*) testing. (Please note that Table 12 now lists the maximum and minimum results for this season and for all past sampling seasons.) *E. coli* is a normal bacterium found in the large intestine of humans and other warm-blooded animals. *E.coli* is used as an indicator organism because it is easily cultured and its presence in the water, in defined amounts, indicates that sewage **MAY** be present. If sewage is present in the water, potentially harmful disease-causing organisms **MAY** also be present.

The *E. coli* concentration was **low (less than 10 counts per 100 mL of sample)** at the **Boat Landing**.

If you are concerned about bacteria levels in this area, you may want to repeat this test next season on a weekend during heavy beach use or after a rain event. Since *E.coli* die quickly in cool pond waters, testing is most accurate and most representative of the health risk to bathers when the source (humans, animals, or waterfowl) is present.



➤ **Table 14: Current Year Biological and Chemical Raw Data**

This table lists the most current sampling season results. Since the maximum, minimum, and annual mean values for each parameter are not shown on this table, this table displays the current year “raw” (meaning unprocessed) data. The results are sorted by station, depth zone (epilimnion, metalimnion, and hypolimnion) and parameter.

➤ **Table 15: Station Table**

As of the Spring of 2004, all historical and current year VLAP data are included in the DES Environmental Monitoring Database (EMD). To facilitate the transfer of VLAP data into the EMD, a new station identification system had to be developed. While volunteer monitoring groups can still use the sampling station names that they have used in the past (and are most familiar with), an EMD station name also exists for each VLAP sampling location. For each station sampled at your lake or pond, Table 15 identifies what EMD station name corresponds to the station names you have used in the past and will continue to use in the future.

### **DATA QUALITY ASSURANCE AND CONTROL**

#### **Annual Assessment Audit:**

During the annual visit to your pond, the biologist conducted a “Sampling Procedures Assessment Audit” for your monitoring group. Specifically, the biologist observed the performance of your monitoring group while sampling and filled out an assessment audit sheet to document the ability of the volunteer monitors to follow the proper field sampling procedures (as outlined in the VLAP Monitor’s Field Manual). This assessment is used to identify any aspects of sample collection in which volunteer monitors fail to follow proper procedures, and also provides an opportunity for the biologist to retrain the volunteer monitors as necessary. This will ultimately ensure that the samples that the volunteer monitors collect are truly representative of actual lake and tributary conditions.

Overall, your monitoring group did an **excellent** job collecting samples on the annual biologist visit this season! Specifically, the members of your monitoring group followed the proper field sampling procedures and there was no need for the biologist to provide additional training. Keep up the good work!

**USEFUL RESOURCES**

*Acid Deposition Impacting New Hampshire's Ecosystems*, NHDES Fact Sheet ARD-32, (603) 271-2975 or [www.des.state.nh.us/factsheets/ard/ard-32.htm](http://www.des.state.nh.us/factsheets/ard/ard-32.htm).

*Best Management Practices to Control Nonpoint Source Pollution: A Guide for Citizens and Town Officials*, NHDES Booklet WD-03-42, (603) 271-2975.

*Canada Geese Facts and Management Options*, NHDES Fact Sheet BB-53, (603) 271-2975 or [www.des.state.nh.us/factsheets/bb/bb-53.htm](http://www.des.state.nh.us/factsheets/bb/bb-53.htm).

*Erosion Control for Construction in the Protected Shoreland Buffer Zone*, NHDES Fact Sheet WD-SP-1, (603) 271-2975 or [www.des.state.nh.us/factsheets/sp/sp-1.htm](http://www.des.state.nh.us/factsheets/sp/sp-1.htm).

*Impacts of Development Upon Stormwater Runoff*, NHDES Fact Sheet WD-WQE-7, (603) 271-2975 or [www.des.state.nh.us/factsheets/wqe/wqe-7.htm](http://www.des.state.nh.us/factsheets/wqe/wqe-7.htm).

*Lake Protection Tips: Some Do's and Don'ts for Maintaining Healthy Lakes*, NHDES Fact Sheet WD-BB-9, (603) 271-2975 or [www.des.state.nh.us/factsheets/bb/bb-9.htm](http://www.des.state.nh.us/factsheets/bb/bb-9.htm).

*Low Impact Development Hydrologic Analysis*. Manual prepared by Prince George's County, Maryland, Department of Environmental Resources. July 1999. To access this document, visit [www.epa.gov/owow/nps/lid\\_hydr.pdf](http://www.epa.gov/owow/nps/lid_hydr.pdf) or call the EPA Water Resource Center at (202) 566-1736.

*Low Impact Development: Taking Steps to Protect New Hampshire's Surface Waters* NHDES Fact Sheet WD-WMB-16, (603) 271-2975 or [www.des.state.nh.us/factsheets/wmb/wmb-17.htm](http://www.des.state.nh.us/factsheets/wmb/wmb-17.htm).

*Proper Lawn Care In the Protected Shoreland, The Comprehensive Shoreland Protection Act*, NHDES Fact Sheet WD-SP-2, (603) 271-2975 or [www.des.state.nh.us/factsheets/sp/sp-2.htm](http://www.des.state.nh.us/factsheets/sp/sp-2.htm).

*Sand Dumping - Beach Construction*, NHDES Fact Sheet WD-BB-15, (603) 271-2975 or [www.des.state.nh.us/factsheets/bb/bb-15.htm](http://www.des.state.nh.us/factsheets/bb/bb-15.htm).

*Soil Erosion and Sediment Control on Construction Sites*, NHDES Fact Sheet WQE-6, (603) 271-2975 or [www.des.state.nh.us/factsheets/wqe/wqe-6.htm](http://www.des.state.nh.us/factsheets/wqe/wqe-6.htm).

*Watershed Districts and Ordinances*, NHDES Fact Sheet WD-WMB-16, (603) 271-2975 or [www.des.state.nh.us/factsheets/wmb/wmb-16.htm](http://www.des.state.nh.us/factsheets/wmb/wmb-16.htm).